

Practitioner's Docket No. 700953-45394C

PATENT

Preliminary Classification:

Proposed Class:

Subclass:

NOTE: "All applicants are requested to include a preliminary classification on newly filed patent applications. The preliminary classification, preferably class and subclass designations, should be identified in the upper right-hand corner of the letter of transmittal accompanying the application papers, for example 'Proposed Class 2, subclass 129.'" M.P.E.P. Section 601, 7th ed.

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Box Patent Application  
Assistant Commissioner for Patents  
Washington, D.C. 20231

## NEW APPLICATION TRANSMITTAL

Transmitted herewith for filing is the patent application of  
Inventor(s): Jeffrey, SCHLOM; Dennis, PANICALI

## CERTIFICATION UNDER 37 C.F.R. SECTIONS 1.8(a) AND 1.10\*

(When using Express Mail, the Express Mail label number is **mandatory**;  
Express Mail certification is optional.)

I hereby certify that, on the date shown below, this correspondence is being:

## MAILING

- ☐ deposited with the United States Postal Service in an envelope addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231.  
37 C.F.R. Section 1.8(a) 37 C.F.R. Section 1.10\*

- ☐ with sufficient postage as first class mail.

- ☒ as "Express Mail Post Office to Address"  
Mailing Label No. \_\_\_\_\_ (mandatory)

EL565095244US

## TRANSMISSION

- ☐ transmitted by facsimile to the Patent and Trademark Office (703) \_\_\_\_\_

Date: October 20, 2000

Signature

Patricia Turner

(type or print name of person certifying)

**\*WARNING:** Each paper or fee filed by "Express Mail" **must** have the number of the "Express Mail" mailing label placed thereon prior to mailing. 37 C.F.R. Section 1.10(b).  
"Since the filing of correspondence under [Section] 1.10 without the Express Mail mailing label thereon is an oversight that can be avoided by the exercise of reasonable care, requests for waiver of this requirement will **not** be granted on petition." Notice of Oct. 24, 1996, 60 Fed. Reg. 56,439, at 56,442.

For (title):GENERATION OF IMMUNE RESPONSES TO PROSTATE-SPECIFIC ANTIGEN (PSA)

**1. Type of Application**

This new application is for a(n)

- ☐ Original (nonprovisional)
- ☐ Design
- ☐ Plant
- ☐ Divisional.
- ☒ Continuation.
- ☐ Continuation-in-part (C-I-P).

**2. Benefit of Prior U.S. Application(s) (35 U.S.C. Sections 119(e), 120, or 121)**

- ☒ The new application being transmitted claims the benefit of prior U.S. application(s). Enclosed are ADDED PAGES FOR NEW APPLICATION TRANSMITTAL WHERE BENEFIT OF PRIOR U.S. APPLICATION(S) CLAIMED.

**3. Papers Enclosed**

**A. Required for Filing Date under 37 C.F.R. Section 1.53(b) (Regular) or 37 C.F.R. Section 1.153 (Design) Application**

44 Pages of Specification  
3 Pages of Claims  
2 Sheets of Drawing

**B. Other Papers Enclosed**

1 Pages of declaration and power of attorney  
1 Pages of Abstract  
       Other

**4. Additional Papers Enclosed**

- ☐ Amendment to claims
  - ☐ Cancel in this applications claims \_\_\_\_\_ before calculating the filing fee. (At least one original independent claim must be retained for filing purposes.)
  - ☐ Add the claims shown on the attached amendment. (Claims added have been

numbered consecutively following the highest numbered original claims.)

- ☐ Preliminary Amendment
- ☐ Information Disclosure Statement (37 C.F.R. Section 1.98)
- ☐ Form PTO-1449 (PTO/SB/08A and 08B)
- ☐ Citations
- ☐ Declaration of Biological Deposit
- ☐ Submission of "Sequence Listing," computer readable copy and/or amendment pertaining thereto for biotechnology invention containing nucleotide and/or amino acid sequence.
- ☐ Authorization of Attorney(s) to Accept and Follow Instructions from Representative
- ☐ Special Comments
- ☐ Other

**5. Declaration or Oath (including power of attorney)**

- ☒ Enclosed

Executed by

- ☒ inventor(s).
- ☐ legal representative of inventor(s). 37 C.F.R. Section 1.42 or 1.43.
- ☐ joint inventor or person showing a proprietary interest on behalf of inventor who refused to sign or cannot be reached.

- ☐ This is the petition required by 37 C.F.R. Section 1.47 and the statement required by 37 C.F.R. Section 1.47 is also attached. See item 13 below for fee.

- ☐ Not Enclosed.

- ☐ Application is made by a person authorized under 37 C.F.R. 1.41 on behalf of *all* the above named inventor(s).

- ☐ Showing that the filing is authorized.  
(*not required unless called into question. 37 C.F.R. Section 1.41(d)*)

**6. Inventorship Statement**

The inventorship for all the claims in this application are:

- ☐ The same.

**or**

- ☐ Not the same. An explanation, including the ownership of the various claims at the time the last claimed invention was made,
  - ☐ is submitted.
  - ☐ will be submitted.

**7. Language**

- ☒ English  
☐ Non-English

- ☐ The attached translation includes a statement that the translation is accurate. 37 C.F.R. Section 1.52(d).

**8. Assignment**

- ☐ An assignment of the invention to \_\_\_\_\_  
\_\_\_\_\_

- ☐ is attached. A separate ☐ "COVER SHEET FOR ASSIGNMENT (DOCUMENT) ACCOMPANYING NEW PATENT APPLICATION" or ☐ FORM PTO 1595 is also attached.

- ☐ will follow.

**9. Certified Copy**

Certified copy(ies) of application(s)

Country	Appln. no.	Filed
Country	Appln. no.	Filed
Country	Appln. no.	Filed

from which priority is claimed

- ☐ is (are) attached.  
☐ will follow.

**10. Fee Calculation (37 C.F.R. Section 1.16)**

- A. ☒ Regular application

\_\_\_\_\_  
CLAIMS AS FILED  
\_\_\_\_\_

Claims	Number Filed	Basic Fee Allowance	Number Extra	Rate	Basic Fee 37 C.F.R. Section 1.16(a) \$710.00
--------	--------------	---------------------	--------------	------	---

Total Claims (37 C.F.R. Section 1.16(c))		18 - 20 =	x	\$18.00	
---	--	-----------	---	---------	--

---

Independent Claims (37 C.F.R. Section 1.16(b))		4 - 3 =	x	\$78.00	\$78.00
---	--	---------	---	---------	---------

---

Multiple Dependent Claim(s), if any (37 C.F.R. Section 1.16(d))					
		+		\$260.00	

---

- ☐ Amendment cancelling extra claims is enclosed.  
☐ Amendment deleting multiple-dependencies is enclosed.  
☐ Fee for extra claims is not being paid at this time.

Filing Fee Calculation \$ 788.00

B. ☐ Design application  
 (\$310.00--37 C.F.R. Section 1.16(f))  
 Filing Fee Calculation \$                     

C. ☐ Plant application  
 (\$480.00--37 C.F.R. Section 1.16(g))  
 Filing Fee Calculation \$                     

# 11. Small Entity Statement(s)

- ☐ Statement(s) that this is a filing by a small entity under 37 C.F.R. Section 1.9 and 1.27 is (are) attached.
- ☐ Status as a small entity was claimed in prior application  
                     /                     , filed on                      from which benefit is being claimed for this application under:

[illegible]

[ ] A copy of the statement in the prior application is included.

## 12. Request for International-Type Search (37 C.F.R. Section 1.104(d))

**13. Fee Payment Being Made at This Time**

[ ] No filing fee is to be paid at this time.  
(This and the surcharge required by 37 C.F.R. Section 1.16(e) can be paid subsequently.)

[ X] Filing fee \$ 788.00

[ ] Petition fee for filing by other than all the inventors or person on behalf of the inventor where inventor refused to sign or cannot be reached  
(\$130.00; 37 C.F.R. Sections 1.47 and 1.17(i))\$

(New Application Transmittal--page 6 of 13)

☐ Processing and retention fee  
(\$130.00; 37 C.F.R. Sections 1.53(d) and 1.21(l))\$

☐ Fee for international-type search report  
(\$40.00; 37 C.F.R. Section 1.21(e)) \$

Total Fees Enclosed \$ 788.00

**14. Method of Payment of Fees**

☒ Check in the amount of \$ 788.00.

☐ Charge Account No. \_\_\_\_\_ in the amount of \$ \_\_\_\_\_.  
A duplicate of this transmittal is attached.

**15. Authorization to Charge Additional Fees**

☒ The Commissioner is hereby authorized to charge the following additional fees by this paper and during the entire pendency of this application to Account No. 50-0850.

☒ 37 C.F.R. Section 1.16(a), (f) or (g) (filing fees)

☐ 37 C.F.R. Section 1.16(b), (c) and (d) (presentation of extra claims)

☐ 37 C.F.R. Section 1.16(e) (surcharge for filing the basic filing fee and/or declaration on a date later than the filing date of the application)

☐ 37 C.F.R. Section 1.17(a)(1)-(5) (extension fees pursuant to Section 1.136(a).

☐ 37 C.F.R. Section 1.17 (application processing fees)

☐ 37 C.F.R. Section 1.18 (issue fee at or before mailing of Notice of Allowance, pursuant to 37 C.F.R. Section 1.311(b))

**16. Instructions as to Overpayment**

☒ Credit Account No. 50-0850.

☐ Refund

  
\_\_\_\_\_  
**SIGNATURE OF PRACTITIONER**

Reg. No. 34,235

David S. Resnick

(type or print name of practitioner)

Tel. No.: (617) 345-6057

101 Federal Street

P.O. Address

Customer No.:

Boston, MA 02110

☒ **Incorporation by reference of added pages**

*(check the following item if the application in this transmittal claims the benefit of prior U.S. application(s) (including an international application entering the U.S. stage as a continuation, divisional or C-I-P application) and complete and attach the ADDED PAGES FOR NEW APPLICATION TRANSMITTAL WHERE BENEFIT OF PRIOR U.S. APPLICATION(S) CLAIMED)*

☒ Plus Added Pages for New Application Transmittal Where Benefit of Prior U.S. Application(s) Claimed

Number of pages added 1

☐ Plus Added Pages for Papers Referred to in Item 4 Above

Number of pages added \_\_\_\_\_

☐ Plus added pages deleting names of inventor(s) named on prior application(s) who is/are no longer inventor(s) of the subject matter claimed in this application.

Number of pages added \_\_\_\_\_

☐ Plus "Assignment Cover Letter Accompanying New Application"

Number of pages added \_\_\_\_\_



☐ **Statement Where No Further Pages Added**

*(if no further pages form a part of this Transmittal, then end this Transmittal with this page and check the following item)*

☐ This transmittal ends with this page.

ADDED PAGE(S) FOR SPECIAL COMMENTS FOR APPLICATION TRANSMITTAL WHERE  
BENEFIT OF PRIOR U.S. APPLICATION(S) IS CLAIMED

Jc813 U.S. PTO  
09/693121  
10/20/00

17. Relate back

A. 35 U.S.C. 119(e)

B. 35 U.S.C. 120, 121 and 365(c)

☒ "This application is a

☒ continuation

☐ continuation-in-part

☐ divisional

of copending application(s)

☒ application number 08/500,306 filed on July 10, 1995.

☐ International Application \_\_\_\_\_ filed on \_\_\_\_\_ and which designated  
U.S."

☐ "The non-provisional application designed above, namely application number 08/500,306  
filed on July 10, 1995, claims benefit of U.S. Provisional Application(s) No(s).:

APPLICATION NO(S).:

FILING DATE

\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

☐ Where more than one reference is made above please combine all references into one sentence.

Added page 1

(Added Page(s) for Special Comments for New Application Transmittal)

Practitioner's Docket No. 953-45394-CPA-2C

PATENT



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of: SCHLOM, et al.

Application No.:  
Filed: Submitted herewith

Group No.:  
Examiner:

For: GENERATION OF IMMUNE RESPONSES TO PROSTATE-SPECIFIC ANTIGEN (PSA)

Assistant Commissioner for Patents  
Washington, D.C. 20231

EXPRESS MAIL CERTIFICATE


"Express Mail" label number EL565095244US  
Date of Deposit 10/20/2000

I hereby state that the following *attached* papers or fees

New Application Transmittal  
Copy of the Application No. 08/500,306 including 47 pages of Specification, Abstract, 2 pages of  
Sequence Listing, 2 sheets of Drawings;  
Copy of Declaration and Power of Attorney;  
Check in the sum of \$788.00; and  
Return Receipt Postcard and Change of Attorney's Address in Application.

is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service  
under 37 C.F.R. section 1.10, on the date indicated above and is addressed to the Assistant Commissioner  
for Patents, Washington, D.C. 20231.

Patricia Turner

  
Signature of person mailing paper or fee

## **GENERATION OF IMMUNE RESPONSES TO PROSTATE-SPECIFIC ANTIGEN (PSA)**

### **FIELD OF THE INVENTION**

5 The present invention relates generally to generation of cellular and humoral immune responses to a mammalian prostate-specific antigen (PSA).

### **BACKGROUND OF THE INVENTION**

10 Cancer of the prostate is the most commonly diagnosed cancer in men and is the second most common cause of cancer death (Carter, et al., 1990; Armbruster, et al., 1993). If detected at an early stage, prostate cancer is potentially curable. However, a majority of cases are diagnosed at later stages when metastasis of the primary tumor has already occurred (Wang, et al., 1982). Even early diagnosis is problematic because not all  
15 individuals who test positive in these screens develop cancer. Present treatment for prostate cancer includes radical prostatectomy, radiation therapy, or hormonal therapy. No systemic therapy has clearly improved survival in cases of hormone refractory disease. With surgical intervention, complete eradication of the tumor is not always achieved and the observed re-occurrence of the cancer (12-68%) is dependent upon the initial clinical  
20 tumor stage (Zietman, et al., 1993). Thus, alternative methods of treatment including prophylaxis or prevention are desirable.

25 Prostate specific antigen (PSA) is a 240 amino acid member of the glandular kallikrein gene family. (Wang, et al., 1982; Wang, et al., 1979; Bilhartz, et al., 1991). PSA is a serine protease, produced by normal prostatic tissue, and secreted exclusively by the epithelial cells lining prostatic acini and ducts (Wang, et al., 1982; Wang, et al., 1979; Lilja, et al., 1993). Prostate specific antigen can be detected at low levels in  
30 the sera of healthy males without clinical evidence of prostate cancer.

However, during neoplastic states, circulating levels of this antigen increase dramatically, correlating with the clinical stage of the disease (Schellhammer, et al., 1993; Huang, et al., 1993; Kleer, et al., 1993; Oesterling, et al., 1991). Prostate specific antigen is now the most widely used marker for prostate cancer. The tissue specificity of this antigen makes PSA a potential target antigen for active specific immunotherapy (Armbruster, et al., 1993; Brawer, et al., 1989), especially in patients who have undergone a radical prostatectomy in which the only PSA expressing tissue in the body should be in metastatic deposits. Recent studies using in-vitro immunization have shown the generation of CD4 and CD8 cells specific for PSA (Peace et al., 1994; Correale et al., 1995). However, although weak natural killer cell responses have been occasionally documented in prostate cancer patients (Choe, et al., 1987), attempts to generate an *in vivo* immune response have met with limited success. For example, several attempts to actively immunize patients with prostate adenocarcinoma cells admixed with Bacillus Calmette-Guerin (BCG) have shown little or no therapeutic benefit (Donovan, et al., 1990). The ability to elicit an immune response as a result of exposure to PSA in vivo would be extremely useful.

Vaccinia virus has been used in the world-wide eradication of smallpox. This virus has been shown to express a wide range of inserted genes, including several tumor associated genes such as p97, HER-2/neu, p53 and ETA (Paoletti, et al., 1993). Other pox viruses that have been suggested as useful for expression of multiple genes include avipox such as fowl pox. Cytokines expressed by recombinant vaccinia virus include IL-1, IL-2, IL-5, IL-6, TNF- $\alpha$  and IFN- $\gamma$  (Paoletti, et al., 1993). Recombinant pox viruses, for example vaccinia viruses, are being considered for use in therapy of cancer because it has been shown in animal models that the co-presentation of a weak immunogen with the highly immunogenic poxvirus

proteins can elicit a strong immune response against the inserted gene product (Kaufman, et al., 1991; Paoletti, et al., 1993; Kantor, et al., 1992a; Kantor, et al., 1992b; Irvine, et al., 1993; Moss, et al., 1993). A recombinant vaccinia virus containing the human carcinoembryonic antigen gene has just completed phase 1 clinical trials in carcinoma patients with no evidence of toxicity other than that observed with the wild type smallpox vaccine (Kantor, et al., 1992b).

Currently, models for the evaluation of prostate therapeutics include the canine (McEntee, et al., 1987), and the Dunning rat (Isaacs, et al., 1986); neither of these models, however, are practical for the study of PSA-recombinant vaccines due to the very low homology of rat and canine PSA to human PSA (Karr, et al., 1995; Schroder, et al., 1982). In contrast, the prostate gland of the rhesus monkey is structurally and functionally similar to the human prostate (Wakui, et al., 1992). At the molecular level, there is 94% homology between either the amino acid or nucleic acid sequences of rhesus PSA (Gauthier, et al., 1993) and those sequences of human prostate specific antigen (Karr, et al., 1995; Lundwall, et al., 1987). Thus, human PSA is essentially an autoantigen in the rhesus monkey. Accordingly, the rhesus monkey can serve as a model for autologous anti-PSA immune reactions.

### **SUMMARY OF THE INVENTION**

We have discovered that by using a recombinant viral vector, preferably a pox virus vector having at least one insertion site containing a DNA segment encoding prostate-specific antigen (PSA), or a cytotoxic T-cell eliciting epitope thereof, operably linked to a promoter capable of expression in the host, a specific humoral and cellular immune response to PSA can be generated. The method preferably comprises introducing a sufficient amount of the recombinant pox virus vector into a host to

stimulate the immune response, and contacting the host with additional PSA at periodic intervals thereafter. The additional PSA, or a cytotoxic T-cell eliciting epitope thereof, may be added by using a second pox virus vector from a different pox genus. In another embodiment, additional PSA can be added by contacting the host with PSA by a variety of other methods, including in one preferred embodiment adding PSA. The PSA may be formulated with an adjuvant or in a liposomal formulation.

In a further embodiment, an immune response to PSA can be generated by contacting the host initially with a sufficient amount of PSA, or a cytotoxic T-cell eliciting epitope thereof, to stimulate an immune response and at periodic intervals thereafter contacting the host with additional PSA. The additional PSA, or a cytotoxic T-cell generating fragment thereof, may be added using a pox virus vector as discussed above.

We have also discovered that human cytotoxic T-cells specific for PSA can be produced using a cytotoxic T-cell eliciting epitope of the PSA and that these cells have the ability to lyse PSA-expressing human prostate carcinoma cells.

As used herein the term "prostate specific antigen" includes the native protein whether purified from a native source or made by recombinant technology, as well as any polypeptide, mutein or portion derived therefrom that is capable of generating an immune response to a native conformationally correct PSA. For example, one can make conservative amino acid substitutions in the molecule without adversely affecting the ability to use the recombinant to generate an antibody that will also recognize native PSA.

5 The pox virus is preferably selected from the group of pox viruses consisting of suipox, avipox, capripox and orthopox virus. Preferred orthopox include vaccinia, rabbit pox and raccoon pox. Preferred avipox includes fowlpox, canary pox and pigeon pox. A more preferred avipox is fowlpox. The preferred suipox is swinepox.

10 Vaccinia viral vectors may elicit a strong antibody response. Thus while numerous boosts with vaccinia vectors are possible, its repeated use may not be preferred in certain instances. We have discovered that by using pox from different genera to boost, this sensitivity problem can be minimized. In accordance with the present invention, in order to avoid such problems, preferably, when the first or initial pox virus vector is vaccinia, the second and subsequent pox virus vectors are selected from the pox viruses from a different genus such as suipox, avipox, capripox or an orthopox immunogenically distinct from vaccinia.

15 Adjuvants include, for example, RIBI Detox, QS21, and incomplete Freund's adjuvant. Liposomal formulations can also be used.

20 Human cytotoxic T-cells specific for PSA produced in accordance with the present invention can be isolated from a human host. These cells can be used in drug assays, used to map cytotoxic T-cells eliciting antigen epitopes or in adoptive cell therapy.

## 25 BRIEF DESCRIPTION OF THE DRAWINGS

30 Figure 1 shows a Western blot of PSA from rV-PSA infected BSC-40 cells. Lanes 2-4 are extracts from supernatant fluid from cells infected overnight with rV-PSA at an MOI of 1, while Lanes 7-9 are extracts from the corresponding infected cells. Lanes 1 and 7 are supernatant extracts and cell extracts from V-Wyeth infected cells. Blot was developed using a



specific MAb for human PSA. This blot illustrates that cells infected with rV-PSA authentically express and secrete the 33 kD PSA protein.

Figures 2A, 2B and 2C show the manifestation of rV-PSA immunization. In Figure 2A, the area of lesions was measured 7 days following each inoculation of rhesus monkeys with either V-Wyeth (open circles) or rV-PSA (closed circles). In Figure 2B, the duration of the lesion was monitored as time of scab disappearance. In Figure 2C, the extent of lymph node swelling was recorded and characterized as very swollen (3+), i.e., more than two axillary nodes swollen; swollen (2+), i.e., one or two nodes easily palpable; marginally swollen (1+), i.e., one node was barely palpable; or not swollen (0), 7 days following inoculation with vaccinia virus. Each symbol represents one monkey.

#### DETAILED DESCRIPTION OF THE INVENTION

We have induced an immune response specific to PSA in the rhesus monkey model by placing the PSA gene into a recombinant viral vector, i.e., a pox vector such as vaccinia virus.

Additionally, an immune response to PSA can be generated by contacting the host initially with a sufficient amount of PSA, or a cytotoxic T-cell eliciting epitope thereof, to stimulate an immune response and at periodic intervals thereafter contacting the host with additional PSA. The additional PSA, or a cytotoxic T-cell generating fragment thereof, may be added using a pox virus vector.

A DNA fragment encoding the open reading frame of human PSA can be obtained, for example, from total RNA extracted from the human metastatic prostate adenocarcinoma cell line, LNCaP.FGC (CRL 1740, American Type Cell Culture (ATCC), Rockville, MD) by reverse

transcriptase PCR using PSA specific oligonucleotide primers 5'  
TCTAGAAGCCCCAAGCTTACCACCTGCA 3' (SEQ. ID. NO.:1), 5'  
TCTAGATCAGGGGTTGGCCACGATGGTGTCTTGATCCACT 3' (SEQ. ID.  
NO.:2). The nucleotide sequence of the PSA cDNA has been published  
(Lundwall, et al., 1987).

Recombinant human PSA can be obtaining using a baculovirus  
expression system in accordance with the method of Bei et al., *J. Clin.  
Lab. Anal.*, 9:261-268 (1995), the disclosure of which is herein  
incorporated by reference.

### **Viral vector**

Basic techniques for preparing recombinant DNA viruses containing a  
heterologous DNA sequence encoding the carcinoma self-associated  
antigen or cytotoxic T-cell eliciting epitope are known to the skilled artisan  
and involve, for example, homologous recombination between the viral  
DNA sequences flanking the DNA sequence in a donor plasmid and  
homologous sequences present in the parental virus (Mackett, et al., *Proc.  
Natl. Acad. Sci. USA* 79:7415-7419 (1982)). For example, recombinant  
viral vectors such as a pox viral vector can be used in delivering the gene.  
The vector can be constructed for example by steps known in the art, e.g.  
analogous to the methods for creating synthetic recombinants of the  
fowlpox virus described in U.S. Patent No. 5,093,258, the disclosure of  
which is incorporated herein by reference. Other techniques include using  
a unique restriction endonuclease site that is naturally present or artificially  
inserted in the parental viral vector to insert the heterologous DNA.

Pox viruses useful in practicing the present invention include  
orthopox, suipox, avipox and capripox virus.

Orthopox include vaccinia, ectromelia and raccoon pox. The preferred orthopox is vaccinia.

Avipox includes fowlpox, canary pox and pigeon pox. The preferred avipox is fowlpox.

Capripox include goatpox and sheeppox.

A preferred suipox is swinepox.

Other viral vectors that can be used include other DNA viruses such as herpes virus and adenoviruses, and RNA viruses such as retroviruses and polio.

For example, the DNA gene sequence to be inserted into the virus can be placed into a donor plasmid, e.g., an *E. coli* plasmid construct, into which DNA homologous to a section of DNA such as that of the insertion site of the poxvirus where the DNA is to be inserted has been inserted. Separately the DNA gene sequence to be inserted is ligated to a promoter. The promoter-gene linkage is positioned in the plasmid construct so that the promoter-gene linkage is flanked on both ends by DNA homologous to a DNA sequence flanking a region of pox DNA which is the desired insertion region. With a parental pox viral vector, a pox promoter is used. The resulting plasmid construct is then amplified by growth within *E. coli* bacteria and isolated. Preferably, the plasmid also contains an origin of replication such as the *E. coli* origin of replication, and a marker such as an antibiotic resistance gene for selection and propagation in *E. coli*.

Second, the isolated plasmid containing the DNA gene sequence to be inserted is transfected into a cell culture, e.g., chick embryo fibroblasts,

along with the parental virus, e.g., poxvirus. Recombination between homologous pox DNA in the plasmid and the viral genome respectively results in a recombinant poxvirus modified by the presence of the promoter-gene construct in its genome, at a site which does not affect virus viability.

As noted above, the gene is inserted into a region (insertion region), in the virus which does not affect virus viability of the resultant recombinant virus. The skilled artisan can readily identify such regions in a virus by, for example, randomly testing segments of virus DNA for regions that allow recombinant formation without seriously affecting virus viability of the recombinant. One region that can readily be used and is present in many viruses is the thymidine kinase (TK) gene. For example, the TK gene has been found in all pox virus genomes examined [leporipoxvirus: Upton, et al., *J. Virology*, 60:920 (1986) (sheep fibroma virus); capripoxvirus: Gershon, et al., *J. Gen. Virol.*, 70:525 (1989) (Kenya sheep-1); orthopoxvirus: Weir, et al., *J. Virol.*, 46:530 (1983) (vaccinia); Esposito, et al., *Virology*, 135:561 (1984) (monkeypox and variola virus); Hruby, et al., *PNAS*, 80:3411 (1983) (vaccinia); Kilpatrick, et al., *Virology*, 143:399 (1985) (Yaba monkey tumor virus); avipoxvirus: Binns, et al., *J. Gen. Virol.* 69:1275 (1988) (fowlpox); Boyle, et al., *Virology*, 156:355 (1987) (fowlpox); Schnitzlein, et al., *J. Virological Methods*, 20:341 (1988) (fowlpox, quailpox); entomopox (Lytvyn, et al., *J. Gen. Virol.* 73:3235-3240 (1992)].

In vaccinia, in addition to the TK region, other insertion regions include, for example, the HindIII M fragment.

In fowlpox, in addition to the TK region, other insertion regions include, for example, the BamHI J fragment [Jenkins, et al., *AIDS Research*

and Human Retroviruses 7:991-998 (1991)] the *EcoRI-HindIII* fragment, *EcoRV-HindIII* fragment, *BamHI* fragment and the *HindIII* fragment set forth in EPO Application No. 0 308 220 A1. [Calvert, et al., *J. of Virol.* 67:3069-3076 (1993); Taylor, et al., *Vaccine* 6:497-503 (1988); Spehner, et al., (1990) and Boursnell, et al., *J. of Gen. Virol.* 71:621-628 (1990)].

In swinepox preferred insertion sites include the thymidine kinase gene region.

In addition to the requirement that the gene be inserted into an insertion region, successful expression of the inserted gene by the modified poxvirus requires the presence of a promoter operably linked to the desired gene, i.e., in the proper relationship to the inserted gene. The promoter must be placed so that it is located upstream from the gene to be expressed. Promoters are well known in the art and can readily be selected depending on the host and the cell type you wish to target. For example in poxviruses, pox viral promoters should be used, such as the vaccinia 7.5K, 40K or fowlpox promoters such as FPV C1A. Enhancer elements can also be used in combination to increase the level of expression. Furthermore, the use of inducible promoters, which are also well known in the art, in some embodiments are preferred.

A specific immune response for PSA can be generated by administering between about  $10^5$ - $10^9$  pfu of the recombinant pox virus, constructed as discussed above to a host, more preferably one uses  $10^7$  pfu. The preferred host is a human. At least one interval thereafter, which is preferably one to three months later, the immune response is boosted by administering additional antigen to the host. More preferably there is at least a second "boost" preferably one to three months after the first boost. The antigen may be administered using the same pox virus vector. The

antigen may preferably be administered using a second pox virus vector from a different pox genera, or may be administered directly using, for example, an adjuvant or liposome. Cytokines, e.g., IL-2, IL-6, IL-12 or co-stimulatory molecules, e.g., B7.1, B7.2, may be used as biologic adjuvants and can be administered systemically to the host or co-administered via insertion of the genes encoding the molecules into the recombinant pox vector.

Adjuvants include, for example, RIBI Detox (Ribi Immunochemical), QS21 and incomplete Freund's adjuvant.

### Generation of Cytotoxic T-Cells

Cytotoxic T-cells specific for PSA can be established from peripheral blood mononuclear cells (PBMC) obtained from a host immunized as discussed above. For example, PBMC can be separated by using Lymphocyte Separation Medium gradient (Organon Teknika, Durham, NC, USA) as previously described [Boyum, et al., *Scand J. Clin Lab Invest* 21: 77-80 (1968)]. Washed PBMC are resuspended in a complete medium, for example, RPMI 1640 (GIBCO) supplemented with 10% pool human AB serum (Pel-Freeze Clinical System, Brown Deer, WI, USA), 2mM glutamine, 100 U/ml penicillin and 100  $\mu$ g/ml of streptomycin (GIBCO). PBMC at a concentration of about  $2 \times 10^5$  cells in complete medium in a volume of, for example, 100  $\mu$ l are added into each well of a 96-well flat-bottom assay plate (Costar, Cambridge, MA, USA). The antigen or peptides are added into the cultures in a final concentration of about 50  $\mu$ g/ml and incubated at 37°C in a humidified atmosphere containing 5% CO<sub>2</sub> for 5 days. After removal of peptide containing media, the cultures are provided with fresh human IL-2 (10U/ml) after 5 days and replenished with IL-2 containing medium every 3 days. Primary cultures are restimulated with the same peptide (50  $\mu$ g/ml) on day 16.  $5 \times 10^5$  irradiated (4,000 rad)

autologous PBMC are added in a volume of about 50  $\mu$ l complete medium as antigen-presenting cells (APC). About five days later, the cultures are provided with human IL-2 containing medium as described previously. Cells are restimulated for 5 days at intervals of 16 days.

5

### Epitope mapping

The cytotoxic T-cells of the present invention can be used to determine the epitope of the PSA that elicits a cytotoxic T-cell. For example, one can cut the PSA into numerous peptide fragments. Alternatively, the fragments can be chemically synthesized. Cytotoxic T-cells can then be plated and different fragments added to different wells. Only T-cells which recognize one of the pre-selected peptide fragments as an epitope will continue to expand, thereby permitting ready identification.

10

15

These fragments can then be used to elicit cytotoxic T-cell instead of using the whole protein. Additionally, one can prepare other fragments containing the epitope to enhance its ability to elicit a cytotoxic T-cell response. Modifications to these fragments are well known in the art and include the use of conjugates, specific amino acid residues such as cystines, etc.

20

### Drug Assay

The cytotoxic T-cell can also be used to screen for compounds which enhance the ability of the antigen to create a cytotoxic T-cell response. For example, cytotoxic T-cells can be incubated with a selected epitope, for example, in a microtiter plate. The compound to be tested, e.g. a drug, is then added to the well and the growth of the T-cells is measured. T-cell expansion indicates that the test compound enhances the T-cell response. Such compounds can be further evaluated.

25

30



## Therapy

The cytotoxic T-cell can be cultured to amplify its number and then injected back into the host by a variety of means. Generally, between  $1 \times 10^5$  and  $2 \times 10^{11}$  cytotoxic T-cells per infusion are administered in, for example, one to three infusions of 200 to 250 ml each over a period of 30 to 60 minutes. After the completion of the infusions, the patient may be treated with recombinant interleukin-2 with a dose of 720,000 IU per kilogram of body weight intravenously every eight hours; some doses can be omitted depending on the patient's tolerance for the drug. In addition, after infusion, additional antigen or fragments containing T-cell eliciting epitope(s) may be administered to the patient to further expand the T-cell number. The antigen or epitope may be formulated with an adjuvant and/or may be in a liposomal formulation.

The cytotoxic T-cells can also be modified by introduction of a viral vector containing a DNA encoding TNF and reintroduced into a host in an effort to enhance the anti-tumor activity of the cells. Other cytokines can also be used.

The recombinant vector can be administered using any acceptable route, including, for example, scarification and injection, e.g., intradermal, subcutaneous, intramuscular, intravenous or intraperitoneal.

For parenteral administration, the recombinant vectors will typically be injected in a sterile aqueous or non-aqueous solution, suspension or emulsion in association with a pharmaceutically-acceptable carrier such as physiological saline.

## REFERENCE EXAMPLE 1

### CONSTRUCTION OF VECTORS



## Pox Viruses

A number of pox viruses have been developed as live viral vectors for the expression of heterologous proteins (Cepko et al., *Cell* 37:1053-1062 (1984); Morin et al., *Proc. Natl. Acad. Sci. USA* 84:4626-4630 (1987); Lowe et al., *Proc. Natl. Acad. Sci. USA*, 84:3896-3900 (1987); Panicali & Paoletti, *Proc. Natl. Acad. Sci. USA*, 79:4927-4931 (1982); Mackett et al., *Proc. Natl. Acad. Sci. USA*, 79:7415-7419 (1982)). Representative fowlpox and swinepox virus are available through the ATCC under accession numbers VR-229 and VR-363, respectively.

### DNA Vectors For *In Vivo* Recombination With A Parent Virus

Genes that code for desired carcinoma associated antigens are inserted into the genome of a pox virus in such a manner as to allow them to be expressed by that virus along with the expression of the normal complement of parent virus proteins. This can be accomplished by first constructing a DNA donor vector for *in vivo* recombination with a pox virus.

In general, the DNA donor vector contains the following elements:

- (i) a prokaryotic origin of replication, so that the vector may be amplified in a prokaryotic host;
- (ii) a gene encoding a marker which allows selection of prokaryotic host cells that contain the vector (e.g., a gene encoding antibiotic resistance);
- (iii) at least one gene encoding a desired protein located adjacent to a transcriptional promoter capable of directing the expression of the gene; and

- (iv) DNA sequences homologous to the region of the parent virus genome where the foreign gene(s) will be inserted, flanking the construct of element (iii).

5 Methods for constructing donor plasmids for the introduction of multiple foreign genes into pox virus are described in W091/19803, the techniques of which are incorporated herein by reference. In general, all DNA fragments for construction of the donor vector, including fragments containing transcriptional promoters and fragments containing sequences homologous to the region of the parent virus genome into which foreign genes are to be inserted, can be obtained from genomic DNA or cloned DNA fragments. The donor plasmids can be mono-, di-, or multivalent (i.e., can contain one or more inserted foreign gene sequences).

10 The donor vector preferably contains an additional gene which encodes a marker which will allow identification of recombinant viruses containing inserted foreign DNA. Several types of marker genes can be used to permit the identification and isolation of recombinant viruses. These include genes that encode antibiotic or chemical resistance (e.g., see Spyropoulos et al., *J. Virol.*, 62:1046 (1988); Falkner and Moss., *J. Virol.*, 62:1849 (1988); Franke et al., *Mol. Cell. Biol.*, 5:1918 (1985), as well as genes such as the *E. coli lacZ* gene, that permit identification of recombinant viral plaques by colorimetric assay (Panicali et al., *Gene*, 47:193-199 (1986)).

#### 25 **Integration Of Foreign DNA Sequences Into The Viral Genome And Isolation Of Recombinants**

Homologous recombination between donor plasmid DNA and viral DNA in an infected cell results in the formation of recombinant viruses that incorporate the desired elements. Appropriate host cells for *in vivo* recombination are generally eukaryotic cells that can be infected by the

virus and transfected by the plasmid vector. Examples of such cells suitable for use with a pox virus are chick embryo fibroblasts, HuTK143 (human) cells, and CV-1 and BSC-40 (both monkey kidney) cells. Infection of cells with pox virus and transfection of these cells with plasmid vectors is accomplished by techniques standard in the art (Panicali and Paoletti, U.S. Patent No. 4,603,112, WO89/03429).

Following *in vivo* recombination, recombinant viral progeny can be identified by one of several techniques. For example, if the DNA donor vector is designed to insert foreign genes into the parent virus thymidine kinase (TK) gene, viruses containing integrated DNA will be TK<sup>+</sup> and can be selected on this basis (Mackett et al., *Proc. Natl. Acad. Sci. USA*, 79:7415 (1982)). Alternatively, co-integration of a gene encoding a marker or indicator gene with the foreign gene(s) of interest, as described above, can be used to identify recombinant progeny. One preferred indicator gene is the *E. coli lacZ* gene: recombinant viruses expressing  $\beta$ -galactosidase can be selected using a chromogenic substrate for the enzyme (Panicali et al., *Gene*, 47:193 (1986)).

### **Characterizing The Viral Antigens Expressed By Recombinant Viruses**

Once a recombinant virus has been identified, a variety of methods can be used to assay the expression of the polypeptide encoded by the inserted gene. These methods include black plaque assay (an *in situ* enzyme immunoassay performed on viral plaques), Western blot analysis, radioimmunoprecipitation (RIPA), and enzyme immunoassay (EIA).

## **EXAMPLE I**

### **Generation of PSA Specific Immune Response**

#### **Materials and Methods**

### *Recombinant Vaccinia Virus*

A 786 bp DNA fragment encoding the entire open reading frame of human prostate specific antigen was amplified by reverse transcriptase PCR (GeneAmp RNA PCR Kit, Perkin Elmer, Norwalk, CT) from total RNA extracted from the human metastatic prostate adenocarcinoma cell line, LNCaP.FGC (CRL 1740, American Type Culture Collection (ATCC), Rockville, MD). The predicted amino acid sequence derived from the PSA coding sequence was shown to be nearly identical to the published sequence (Lundwall, et al., 1987), differing only in a change from asparagine to tyrosine at position 220. The PSA DNA fragment, containing the entire coding sequence for PSA, 41 nucleotides of the 5' untranslated region, and 520 nucleotides of the 3' untranslated region, was inserted into the Xba I restriction endonuclease cleavage site of the vaccinia virus transfer vector pT116. The resulting plasmid, designated pT1001, contains the PSA gene under the control of the vaccinia virus 40K promoter (Gritz, et al. 1990) and the *E. coli lacZ* gene under the control of the fowlpox virus C1 promoter (Jenkins, et al., 1991). The foreign genes are flanked by DNA sequences from the Hind III M region of the vaccinia genome. A plaque-purified isolate from the Wyeth (New York City Board of Health) strain of vaccinia was used as the parental virus in the construction of the recombinant vaccinia virus. The generation of recombinant vaccinia virus was accomplished via homologous recombination between vaccinia sequences in the Wyeth vaccinia genome and the corresponding sequences in pT1001 in vaccinia-infected RK<sub>13</sub> cells (CCL 37, ATCC) transfected with pT1001. Recombinant virus was identified using a chromogenic assay, performed on viral plaques *in situ*, that detects expression of the *lacZ* gene product in the presence of halogenated indolyl-beta-D-galactoside (Bluo-gal), as described previously (Panacali, et al., 1986). Appropriate blue recombinant viruses were purified by four rounds of plaque-purification. Virus stocks were prepared

by clarifying infected RK<sub>13</sub> cell lysates followed by centrifugation through a 36% sucrose cushion.

### *Characterization of Recombinant Virus*

#### Southern Analysis of DNA recombination

The recombinant vaccinia genome was analyzed by viral DNA extraction, restriction endonuclease digestion with Hind III, and Southern blotting as previously described (Kaufman *et al.*, 1991).

#### Western Analysis of protein expression

Confluent BSC-40 cells were infected with either parental wild type vaccinia virus (designated V-Wyeth) or recombinant vaccinia-PSA (designated rV-PSA) at an MOI of 1 in Dulbecco's Modified Eagle's Medium containing 2% fetal bovine serum. After an overnight infection, the medium was removed from the cells, and an aliquot was methanol precipitated to assay for the presence of secreted PSA. The infected cells were lysed in hypotonic lysis buffer (150 mM NaCl, 0.05% EDTA, 10 mM KCl, 1 mM PMSF) and then sonicated. Cell lysates and culture media were electrophoresed on an SDS-10% acrylamide gel. The proteins were transblotted to nitrocellulose, and the blot was incubated with a rabbit antibody specific for PSA (P0798, Sigma Chemical Co., St. Louis, MO) for 4 hours at ambient temperature, washed, and then incubated with goat anti-rabbit phosphatase-labeled secondary antibody (AP, Kirkegaard & Perry Laboratories, Gaithersburg, MD) and developed according to the manufacture's instructions.

#### *Generation of B-cell lines*

Monkey autologous B lymphoblastoid cell lines (BLCL) were established by infecting  $1 \times 10^5$  freshly isolated PBMCs in 100 ml of RPMI 1640 supplemented with L-glutamine, gentamicin, and 10% FCS

(Biofluids, Rockville, MD) with 100 ml supernatant from S594 cells (kindly provided by Dr. M. D. Miller, Harvard Medical School, New England Regional Primate Research Center, Southborough, MA), which contains the baboon herpesvirus *Herpes papio*, in a 96 well, flat-bottomed plate (Costar, Cambridge, MA). Following transformation, cells were expanded, and media changed once weekly.

### *Immunization of Monkeys*

Twelve juvenile male rhesus monkeys (*Macaca mulatta*), ages 1 to 2 years, were assigned to three vaccination groups of four animals each. One animal from each group was prostatectomized. Animals were immunized 3 times on days 1, 29, and 57. Doses of either  $1 \times 10^7$  or  $1 \times 10^8$  PFU of rV-PSA were administered to 4 animals by skin scarification. V-Wyeth ( $1 \times 10^8$  PFU) was administered to 4 animals as controls. The animals were housed and maintained at the Toxicology Research Laboratory, University of Illinois at Chicago (TRL/UIC) in accordance with the guidelines of the National Cancer Institute Animal Care and Use Committee and the Guide for the Care and Use of Laboratory Animals (Department of Health and Human Services Publication NIH 85-23, revised 1985 by the FDA Center for Biologics Evaluation and Research Office of Biological Product Review, Division of Product Quality Control, Pathology and Primatology Laboratory, Bethesda, MD).

### *Toxicology*

Physical examinations were performed on ketamine (Ketamine® HCl, 10 mg/kg I.M.) sedated animals. Rectal temperatures and weights were recorded for each monkey on a weekly basis. The vaccination site was observed and erythema and swelling were measured by caliper. Each animal was examined for regional lymphadenopathy, hepatomegaly, and splenomegaly. Any other gross abnormalities were also recorded.

Blood was obtained by venipuncture from the femoral vein of ketamine sedated animals before and after each immunization. A complete blood count, differential, hepatic and renal chemistry evaluation was performed on each monkey by TRL/UIC. Results were compared to normal primate values (Kantor *et al.*, 1992b). Circulating levels of PSA before and after immunization were analyzed by radioimmunoassay (Tandem™, Hybritech, San Diego, CA).

#### *Measurement of Antibody Titers*

Prior to each immunization and 2 weeks following each immunization, anti-PSA antibody was quantified by ELISA. Microtiter plates were coated with purified PSA (100 ng/well, Calbiochem, La Jolla, CA), ovalbumin (100 ng/well, Sigma), or  $1 \times 10^7$  PFU/well UV-inactivated V-Wyeth in PBS. The plates were blocked with 2% BSA in PBS, dried, and stored at  $-20^{\circ}$  C until used. The plates were incubated with serum diluted 1:5, as well as a monoclonal antibody for PSA (DAKO M750, Denmark) as a standard control, for 24 hours at  $4^{\circ}$  C. Plates were washed several times with PBS containing 1% BSA, and incubated at  $37^{\circ}$  C for 45 min with horseradish peroxidase-conjugated goat anti-human IgG or IgM heavy chain specific antiserum (1:8000) (Southern Biotechnology Associates, Birmingham, AL) and antibody detected by HRP substrate system (Kirkegaard & Perry Laboratories, Gaithersburg, MD) according to the manufacture's instructions. The absorbance of each well was read at 405 nm using a Bio-Tek EL310 microplate ELISA reader (Winooski, VT).

#### *Lymphoproliferative Assay*

Autologous monkey BLCL were plated at a density of  $3 \times 10^6$  cells/well in 24 well plates with 160 mg/well purified PSA (Fitzgerald, Concord, MA) or 160 mg/well ovalbumin (Sigma) at  $37^{\circ}$  C for 24 hours. Cells were then  $\gamma$ -irradiated (14000 rad), harvested, washed and



suspended at a final concentration of  $1 \times 10^7$ /ml. Fresh monkey PBMCs from heparinized blood, 6 weeks to 7 months after the last immunization, were isolated on lymphocyte separation medium (Organon Teknika, West Chester, PA). Lymphoproliferative responses were evaluated by co-culturing  $1.5 \times 10^5$  cells with  $5 \times 10^5$  cells/well of autologous BLCL in 0.2 ml of RPMI 1640 supplemented with 10% heat-inactivated fetal calf serum in flat-bottomed 96 well plates (Costar) for 5 days. PBMCs were cultured with  $2 \times 10^7$  PFU/ml UV-inactivated V-Wyeth as a recall antigen or 2 mg/ml Con-A as positive controls. Cells were labeled for the final 12-18 h of the incubation with 1 mCi/well [ $^3$ H]thymidine (New England Nuclear, Wilmington, DE) and harvested with a PHD cell harvester (Cambridge Technology, Cambridge, MA). The incorporated radioactivity was measured by liquid scintillation counting (LS 6000IC; Beckman, Duarte, CA). The results from triplicate wells were averaged and are reported as mean  $\pm$  SEM.

## Results

### *Generation and Characterization of Recombinant Virus*

The cDNA fragment encoding the open reading frame of human PSA was obtained by reverse transcriptase PCR using PSA specific oligonucleotide primers 5' TCTAGAAGCCCCAAGCTTACCACCTGCA 3' (SEQ. ID. NO.:1), 5' TCTAGATCAGGGGTTGGCCACGATGGTGTCTTGATCCACT 3' (SEQ. ID. NO.:2), and ligated into the vaccinia virus transfer vector pT106. This vector contains a strong vaccinia virus early/late promoter (designated P40) upstream of the multiple cloning site to drive the synthesis of the inserted gene product. The ligation and orientation of the PSA DNA fragment, as well as promoter position were verified by PCR and sequencing. The chimeric vector construct was inserted into the vaccinia virus genome Hind III M site by homologous recombination as previously



reported (Kaufman, et al., (1991)), and confirmed by Southern analysis probing with <sup>32</sup>P radiolabeled DNA corresponding to PSA sequences and vaccinia sequences in the Hind III M region (data not shown). The entire cDNA sequence of PSA in the vaccinia virus clone was shown to be nearly identical to the published sequences (Lundwall, et al., 1987).

Expression of recombinant protein was confirmed by western blot analysis of supernatant fluids and protein extracts from rV-PSA infected BSC-40 cells. These cells are routinely used for the evaluation of recombinant vaccinia products (Moss, et al., 1993). Incubation of cell supernatant blots from rV-PSA infected cells with rabbit anti-PSA antibody revealed a single immunoreactive polypeptide of approximately 33,000 daltons (Figure 1, lanes 2-4). Similarly, incubation of protein extract blots from rV-PSA infected cells revealed a single band of the same molecular weight (Figure 1, lanes 7-9). This is consistent with the predicted size of the PSA molecule (Armbruster, et al., 1993; Wang, et al., 1982). Cell supernatant blots (lane 1) or protein extract blots (lane 6) from cells infected with parental strain V-Wyeth remained negative for expression of PSA. These results thus demonstrate that a recombinant vaccinia virus can faithfully express the human PSA gene product.

#### *Rhesus Monkey Model*

The prostate gland of the rhesus monkey is structurally and functionally similar to the human prostate (Wakui, et al., 1992). At the molecular level, there is 94% homology between both the amino acid and nucleic acid sequences of rhesus PSA (Gauthier, et al., 1993) and human prostate specific antigen (Karr, et al., 1995; Lundwall, et al., 1987). Human PSA is essentially an autoantigen in the rhesus monkey.

#### *Experimental Design*

Table 1 delineates the protocol used in the immunization of 12 rhesus monkeys with either rV-PSA or the control V-Wyeth by skin scarification. Three groups of 4 animals were immunized with either rV-PSA at  $1 \times 10^7$  PFU/dose, rV-PSA at  $1 \times 10^8$  PFU/dose, or V-Wyeth at  $10^8$  PFU/dose 3 times at 4 week intervals. These doses were chosen to ascertain the maximum tolerated dose for safety as well as to obtain maximum humoral and cell-mediated responses to PSA.

The rhesus monkeys were divided into 3 groups: high dose V-Wyeth, low dose rV-PSA, and high dose rV-PSA. One animal in each group was surgically prostatectomized to parallel two situations with regard to potential therapy in humans: (a) prostate intact, with primary and/or metastatic disease; or (b) patients prostatectomized with prostate cancer metastatic deposits. The presence of an intact prostate gland could conceivably serve as an antigen 'sink', either inducing anergy through persistence of antigen, or masking immunological effects by sequestering reactive cells or antibodies.

#### *Physical Consequence of Immunization*

The area of the lesions induced by rV-PSA or V-Wyeth was analyzed 7 days following each inoculation. In general, more induration was seen after the first inoculation, compared to the second inoculation (Figure 2A). After the third inoculation, there was no swelling of the vaccination site. The duration of the lesion following each immunization was shorter after each inoculation (Figure 2B). Regional lymph node swelling following vaccination was greater in most monkeys following the first immunization, compared to the second, or third immunization (Figure 2C). In general, no differences were seen in these parameters with the use of rV-PSA or V-Wyeth. Monkeys receiving V-Wyeth were compared with those receiving rV-PSA with respect to constitutional symptoms. Mild temperature

elevations were seen in all animals following vaccination. There was no evidence of weight loss, hepatomegaly or splenomegaly in any of the animals, and there was no differences between V-Wyeth or rV-PSA treated animals (data not shown). Animals were tested for complete blood count, differential, and hepatic and renal chemistries. Complete blood counts remained within normal limits throughout the study in both V-Wyeth and rV-PSA immunized animals (Table 2). Hepatic and renal functions were assessed prior to immunization and 12 weeks following primary immunization (Table 3). Parameters analyzed included alkaline phosphatase, blood urea nitrogen, alanine aminotransferase, aspartate aminotransferase, lactate dehydrogenase, and creatine and creatine kinase levels. There was no significant difference between animals receiving V-Wyeth or rV-PSA. There was no detectable PSA in the circulation of any of these monkeys after any immunization (detection limit was 0.1 ng/ml). At this time, which is 54 weeks post all immunizations, no toxicities were observed in monkeys of any of the groups, including those which were prostatectomized.

#### *PSA Specific Humoral Responses*

As indicated in Table 1, monkeys 1-4 were administered V-Wyeth while monkeys 5-12 were administered rV-PSA. Sera from each of these monkeys were analyzed by ELISA for immunoreactivity to PSA or UV-inactivated V-Wyeth, and ovalbumin as control antigen. Sera obtained from monkeys prior to vaccination were negative for reactivity to PSA (Table 4, PI). Fifteen days following primary immunization, monkeys in both the  $1 \times 10^8$  and  $1 \times 10^7$  dose rV-PSA groups developed low titer IgM antibodies specific for PSA (titers were determined at a 1:5 serum dilution). Although other isotopes of antibody were analyzed (IgG, IgA, IgM), only IgM was induced by rV-PSA throughout the observation period of 270 days. The antibody titers decreased over the 4 weeks prior to the

next inoculation. Prior to the second vaccination on day 29, 3 of 4 animals in the  $1 \times 10^7$  rV-PSA group remained positive for PSA antibody, while 4 of 4 animals remained positive in the  $1 \times 10^8$  rV-PSA group. Anti-PSA antibody titers increased after the second vaccination on day 29, but remained static after the third vaccination on day 57. By 270 days after the primary immunization, all animals were negative for PSA IgM antibody. Monkeys remained negative for IgG specific for PSA throughout the observation period (data not shown). There was no correlation between rV-PSA dose and anti-PSA IgM titer, nor was there any apparent effect of prostatectomy. All monkey sera were negative for IgG or IgM to ovalbumin at all time points; as a positive control, however, the IgG titer in all three treatment groups to vaccinia virus was greater than 1:2000 as early as 29 days after the primary immunization (data not shown).

In general, vaccinia virus is a weak human pathogen (Paoletti *et al.*, 1993). Following vaccination, local erythema, induration, low-grade fever, and regional lymphadenopathy are common. The virus replicates in the epidermal cells of the skin and the virus is usually cleared within 14 days. All monkeys, whether given V-Wyeth or rV-PSA, exhibited the usual low grade constitutional symptoms of a vaccinia virus infection (Figure 2). There was no evidence of any adverse effects as indicated by changes in blood counts, differentials, hepatic and renal chemistries (Tables 2-3). The monkeys appeared healthy, without any physical signs of toxicity, throughout the 54 weeks of observation.

Although the rV-PSA construct was unable to elicit an anti-PSA IgG response, PSA specific IgM responses were noted in all rV-PSA immunized monkeys regardless of dose level (Table 4). These antibody responses were of low titer, short lived and could not be boosted, indicating induction of a primary response but not memory B-cells or affinity maturation.

### *PSA Specific Lymphoproliferative Assay*

PSA specific T-cell responses in monkeys immunized with rV-PSA or V-Wyeth were analyzed using a lymphoproliferative assay. As seen in Table 5, the PBMCs from all monkeys analyzed responded, regardless of whether they received rV-PSA or V-Wyeth, to the lymphocyte mitogen concanavalin-A, as well as with the recall antigen UV-inactivated V-Wyeth. Differential responses to PSA *versus* medium alone or ovalbumin were seen in 1 animal (number 6) in the  $1 \times 10^7$  PFU rV-PSA group. All PBMCs from animals in the  $1 \times 10^8$  PFU rV-PSA group, however, responded to PSA in this assay. This experiment was repeated 5 times with similar results and data shown in Table 5 is from PBMCs isolated from monkeys 270 days after the primary immunization. No differences in PSA specific T-cell responses were seen in the prostatectomized monkeys.

To investigate cell mediated responses to the administration of rV-PSA, lymphoproliferative assays were performed using PBMCs from animals receiving the recombinant vaccine. One of four monkeys receiving the lower dose of rV-PSA ( $1 \times 10^7$  PFU) and four of four receiving the higher dose ( $1 \times 10^8$  PFU) maintained specific T-cell responses to PSA protein up to 270 days following primary immunization as indicated by the lymphoproliferative assay (Table 5). Prostatectomy appeared to have no effect on either the humoral or cellular responses of monkeys receiving rV-PSA. Evidence of PSA specific T-cell responses in monkeys lacking mature antibody isotopes could be due to two distinct events following vaccination with rV-PSA: a T-cell independent event, leading to IgM production, and a T-cell dependent event, leading to specific lymphoproliferative responses.

Table 1 Inoculation protocol of rhesus monkeys with the  
PSA recombinant and wild-type vaccinia virus

Monkey	Prostate	Immunogen	Dose* (PFU)
1	Yes	V-Wyeth	$1 \times 10^8$
2	Yes	V-Wyeth	$1 \times 10^8$
3	Yes	V-Wyeth	$1 \times 10^8$
4	No	V-Wyeth	$1 \times 10^8$
5	Yes	rV-PSA	$1 \times 10^7$
6	Yes	rV-PSA	$1 \times 10^7$
7	Yes	rV-PSA	$1 \times 10^7$
8	No	rV-PSA	$1 \times 10^7$
9	Yes	rV-PSA	$1 \times 10^8$
10	Yes	rV-PSA	$1 \times 10^8$
11	Yes	rV-PSA	$1 \times 10^8$
12	No	rV-PSA	$1 \times 10^8$

\* All animals received 3 immunizations at 4 week intervals.

Table 2  
Mean WBC count, hematocrit, and differential count in rhesus monkeys receiving recombinant or wild-type vaccine

Test	Normal ranges	V-Wyeth (n=4)		rV-PSA (n=8)	
		Before	After	Before	After
		immunization <sup>a</sup>	immunization <sup>b</sup>	immunization	immunization
WBC	7-15 x 10 <sup>3</sup>	5.0 ± 0.8	5.1 ± 0.5	5.2 ± 0.7	5.8 ± 0.9
Hematocrit (vol. %)	33-43	37.4 ± 0.2	37.0 ± 0.1	37.8 ± 0.4	37.0 ± 0.5
Lymphocytes	1-7 x 10 <sup>3</sup>	2.8 ± 0.7	3.9 ± 0.5	2.2 ± 0.4	3.5 ± 0.8
SEGS <sup>c</sup> (%)	3-69	2.0 ± 0.2	0.78 ± 0.2	2.9 ± 0.6	1.9 ± 0.3
Monocytes (%)	0-8	0.1 ± 0.05	0.2 ± 0.04	0.1 ± 0.04	0.2 ± 0.50
Eosinophils (%)	0-8	0.1 ± 0.02	0.2 ± 0.10	0.1 ± 0.03	0.1 ± 0.02

<sup>a</sup> 1 week prior to primary immunization

<sup>b</sup> 12 weeks following primary immunization

<sup>c</sup> Segmented lymphocytes

Table 3 Mean serum chemistry values in rhesus monkeys receiving recombinant or wild-type vaccine

Test	Normal ranges	V-Wyeth (n=4)		rV-PSA (n=8)	
		Before immunization <sup>a</sup>	After immunization <sup>b</sup>	Before immunization	After immunization
ALKP <sup>c</sup> (u/l)	200-800	451 ± 48	610 ± 33	339 ± 74	454 ± 47
BUN <sup>d</sup> (mg/dl)	12-30	19.0 ± 3.0	17.8 ± 0.9	17.1 ± 0.6	20.5 ± 1.0
ALT <sup>e</sup> (u/l)	20-60	25.2 ± 1.9	22.8 ± 1.0	28.9 ± 5.3	25.8 ± 1.6
AST <sup>f</sup> (u/l)	40-80	37.8 ± 2.3	31.8 ± 4.4	37.9 ± 3.6	31.9 ± 2.4
LDH <sup>g</sup> (u/l)	200-500	194 ± 20	212 ± 21	236 ± 41	194 ± 13
Creatine (mg/dl)	0.5-1.0	0.9 ± 0.10	0.8 ± 0.03	0.8 ± 0.05	0.8 ± 0.02
Creatine Kinase (u/l)	500-2000	662 ± 112	466 ± 119	498 ± 120	563 ± 81

<sup>a</sup> 1 week prior to primary immunization

<sup>b</sup> 12 weeks following primary immunization

<sup>c</sup> Alkaline phosphatase

<sup>d</sup> Blood urea nitrogen

<sup>e</sup> Alanine aminotransferase

<sup>f</sup> Aspartate aminotransferase

<sup>g</sup> Lactate dehydrogenase



Table 4 Primate IgM<sup>a</sup> Response to Inoculation with rV-PSA

Monkey	Immunogen	Dose (PFU)	Days Post Immunization <sup>b</sup>						
			PI <sup>d</sup>	15	29 <sup>e</sup>	43	57 <sup>e</sup>	71	270
1	V-Wyeth	1x10 <sup>8</sup>	ND <sup>f</sup>	ND	ND	ND	ND	ND	ND
2	V-Wyeth	1x10 <sup>8</sup>	ND	ND	ND	ND	ND	ND	ND
3	V-Wyeth	1x10 <sup>8</sup>	ND	ND	ND	ND	ND	ND	ND
4 <sup>c</sup>	V-Wyeth	1x10 <sup>8</sup>	ND	ND	ND	ND	ND	ND	ND
5	rV-PSA	1x10 <sup>7</sup>	ND	>40	5	20	>40	>40	ND
6	rV-PSA	1x10 <sup>7</sup>	ND	>40	ND	ND	20	20	ND
7	rV-PSA	1x10 <sup>7</sup>	ND	>40	5	20	20	20	ND
8 <sup>c</sup>	rV-PSA	1x10 <sup>7</sup>	ND	>40	5	10	>40	>40	ND
9	rV-PSA	1x10 <sup>8</sup>	ND	20	5	20	10	10	ND
10	rV-PSA	1x10 <sup>8</sup>	ND	20	5	40	>40	NT <sup>g</sup>	ND
11	rV-PSA	1x10 <sup>8</sup>	ND	>40	>40	>40	>40	>40	ND
12 <sup>c</sup>	rV-PSA	1x10 <sup>8</sup>	ND	>40	20	40	20	20	ND

<sup>a</sup> All monkey seras were negative for IgG to PSA at all time points;

All seras were positive for IgG to vaccinia virus (>1:2000) at day 71.

<sup>b</sup> Monkeys received vaccinations on days 1, 29, and 57. Sera (1:5) was tested by ELISA. Titers were calculated using an O.D. of 0.4.

<sup>c</sup> Animal was prostatectomized.

<sup>d</sup> PI, Pre-immune.

<sup>e</sup> Animals bled before boosting.

<sup>f</sup> ND, not detectable; limit of detection was <1:5 dilution.

<sup>g</sup> NT, not tested.

Table 5 PSA Specific Lymphoproliferative T-cell Responses of Rhesus PBMCs  
270 Days Following Inoculation with rV-PSA

Monkey	Immunogen	Dose (PFU)	Antigen <sup>a</sup>			
			Medium	Con A	Oval	UV-Wyeth PSA <sup>d</sup>
1	V-Wyeth	1x10 <sup>8</sup>	397	65701	376	24785 414
2 <sup>b</sup>	V-Wyeth	1x10 <sup>8</sup>	NT	NT	NT	NT
3	V-Wyeth	1x10 <sup>8</sup>	450	84860	522	18859 413
4 <sup>c</sup>	V-Wyeth	1x10 <sup>8</sup>	532	107840	553	16571 387
5	rV-PSA	1x10 <sup>7</sup>	412	85276	408	6040 539
6	rV-PSA	1x10 <sup>7</sup>	401	96398	404	7776 <b>3,134</b>
7	rV-PSA	1x10 <sup>7</sup>	417	90801	522	10908 434
8 <sup>c</sup>	rV-PSA	1x10 <sup>7</sup>	1069	99216	744	15346 484
9	rV-PSA	1x10 <sup>8</sup>	384	106248	386	14499 <b>10,635</b>
10	rV-PSA	1x10 <sup>8</sup>	432	92263	404	19872 <b>18,561</b>
11	rV-PSA	1x10 <sup>8</sup>	411	94055	1063	5124 <b>16,245</b>
12 <sup>c</sup>	rV-PSA	1x10 <sup>8</sup>	420	124896	392	11944 <b>12,945</b>

<sup>a</sup> Antigen concentrations were: Con a (2 µg/ml); Ovalbumin (100 µg/ml);

UV-Wyeth (2x10<sup>7</sup> pfu/ml); and PSA (100 µg/ml). Each value represents a mean CPM of triplicate samples. Standard deviation never exceeded 10%.

<sup>b</sup> NT, Not Tested. B-cells were not transformed for this animal.

<sup>c</sup> Animal was prostatectomized.

<sup>d</sup> Values in bold are significant when compared to their respective medium control values (p <0.001).

## EXAMPLE II

### Identification Of Potential Prostate Specific Antigen (PSA) Specific T Cell Epitopes

5

10

15

20

25

Since the entire amino acid sequence of human PSA is known and human class 1 HLA A2 consensus motifs have been described, studies were undertaken to identify a series of peptides that would potentially bind class 1 A2 molecules. A2 was chosen since it is the most common HLA class 1 molecule being represented in approximately 50% of North American Caucasians and 34% of African Americans. The peptide sequence of PSA was thus examined for matches to the consensus motifs for HLA A2 binding peptides. Peptides were only selected if their sequence diverged sufficiently from the PSA-related human glandular kallikrein (HGK) gene and pancreatic kallikrein antigen (PKA) sequences.

The amino acid sequence of human PSA was scanned using a predictive algorithm that combines a search for anchor residues with numerical assignments to all residues at all positions. The T2 cell binding assay was then used to determine which peptides bound human HLA A2 molecules. As can be seen in Table 6, PSA peptides 141-150, 154-163 and 146-154 scored positive in this assay (*Nijman, H.W., et al., Eur. J. Immunol.* 23:1215-1219, 1993). Table 7 gives the amino acid sequence of these peptides and compares them to corresponding sequences of HGK and PKA.

**Table 6**  
**PSA peptide binding assay**

Antigen MAb A2, 69

5	None		127.25 <sup>a</sup>
	PSA 141-150	230.34	
	PSA 146-154	223.97	
	PSA 154-163	182.30	
10	Peptides were used at a concentration of 50µg/ml		
	<sup>a</sup> Mean channel fluorescent intensity.		
	CIRA2 cell line was used as positive control for anti-A2 staining [99.4 (241.15)].		

**Table 7**  
**PSA peptide amino acid sequence**

PSA	141-150	F	L	T	P	K	K	L	Q	C	V
HGK		-	-	R	-	R	S	-	-	-	-
PKA		-	S	F	-	D	D	-	-	-	-
PSA	146-154	K	L	Q	C	V	D	L	H	V	
HGK		S	-	-	-	-	S	-	-	L	
PKA		D	-	-	-	-	-	-	K	I	
PSA	154-163	V	I	S	N	D	V	C	A	Q	V
HGK		L	L	-	-	-	M	-	-	R	A
PKA		I	L	P	-	-	E	-	E	K	A

**EXAMPLE III**  
**Establishment of Human T Cell Lines**  
**Cytolytic For Human Tumor Cells Expressing PSA**

PBMC from normal healthy donors expressing the HLA A2 class 1 allele were used in an attempt to determine if PSA specific peptides are immunogenic for humans. Peptides 141-150 and 154-163 were used in this study. The methodology used for the establishment of these cell lines involves pulsing of PBMC with peptide and IL-2 as previously described (*Tsang, K.Y., et al. JNCI, in press* and in U.S. Application Serial No. 08/396,385, the disclosure of which is herein incorporated by reference). T cell lines were able to be established from 5/6 normal donors using PSA peptide 141-150 and from 6/6 normal donors using PSA peptide 154-163. Moreover, PBMC were obtained from two prostate cancer patients. T cell lines were established from these PBMC cultures using peptide 154-163.

Some of these T cell lines have been phenotyped. As seen in Table 8, one cell line designated T-866, which was derived from pulsing with peptide 141-150, contains appreciable amounts of CD4 + /CD8 + double positive cells and another cell line, designated T-1538, derived from pulsing with peptide 154-163, shows a similar phenotype.

Four of the T cell lines derived from three different individuals were then assayed for their ability to lyse human cells (Table 9). As seen in Table 9, the T cell line designated T-866, derived from peptide 141-150, was able to lyse T2 cells when pulsed with the appropriate peptide (141-150). No lysis was seen using the PSA negative human colon cancer cell line COLO-205. While 80% lysis was seen using the LNCAP PSA containing human prostate cancer cell line. When employing the NK target K562, which measures non-specific lysis due to NK cell activity, only 23% lysis was obtained. Similar results were seen employing a different T cell

line obtained from the same patient which was derived from pulsing with PSA peptide 154-163. Two additional T cell lines which were derived from peptide 154-163 were also analyzed. One was from a normal donor (T-1538) and one was from a prostate cancer patient (T-PC2). As can be seen in Table 9, employing both of these T cell lines, enhanced lysis was seen when the T2 cell line was pulsed with the 154-163 peptide and enhanced lysis was seen when employing the PSA expressing prostate specific cell line LNCAP, as compared to COLO-205 or K562. These studies demonstrate that T cell lines can be established using the peptides and protocols generated here which have the ability to lyse PSA expressing human prostate carcinoma cells.

**Table 8**  
**Flow cytometry analysis of PSA peptide specific T cells**

T-cell Line	PSA Peptide	CD3	CD4	CD8	CD4/CD8	CD56
T-866	141-150	96	35	6.5	59	0
T-1538	154-163	94	5.2	32	62	0

Table 9  
Cytotoxic effects of PSA peptide specific T cells

T-cell Line	PSA Peptide	% specific release (lysis)				
		T2	T2 + peptide	LNCAP	K562	COLO-205
T-866	141-150	10 <sup>a</sup>	40	80	23	7
T-866	154-160	16	35	60	22	10
T-1538	154-160	10	40	29	3	10
T-PC2	154-160	15	35	35	2	8

<sup>a</sup>Percent of <sup>111</sup>In specific release  
24 hour cytotoxic assay (E:T ratio, 25:1) (SD < 2.5%)

The following is a listing of publications referred to in the foregoing specification.

- 5 Armbruster, D. A. Prostate-specific antigen: biochemistry, analytical methods, and clinical application. *Clinical Chemistry*, **39**:181-195, (1993).
- 10 Bilhartz, D. L., Tindall, D. J., and Oesterling, J. E. Prostate-specific antigen and prostatic acid phosphatase: biomolecular and physiological characteristics. *Urology*, **38**:95-102, (1991).
- 15 Brawer, M. K., and Lange, P. H. Prostate-specific antigen and premalignant change: implications for early detection. *CA Cancer Journal Clinic*, **39**:361-375, (1989).
- 20 Carter, H. B., and Coffey, D. S. The prostate: an increasing medical problem. *Prostate*, **16**:39-48, (1990).
- 25 Chatterjee, M. B., Foon, K. A., and Kohler, H. Idiotypic antibody immunotherapy of cancer. *Cancer Immunology and Immunotherapy*, **38**:75-82, (1994).
- 30 Cheever, M. A., Chen, W., Disis, M. T., and Peace, D. J. T-cell immunity to oncogenic proteins including mutated RAS and chimeric BCR-ABL. *Annals of the New York Academy of Science*, **690**:101-112, (1993).
- 35 Choe, B. K., Frost, P., Morrison, M. K., and Rose, N R. Natural killer cell activity of prostatic cancer patients. *Cancer Investigations*, **5**:285-291, (1987).
- 40 Conry, R. M., Salch, M. N., Schlom, J., and LoBuglio, A. F. Breaking tolerance to carcinoembryonic antigen with a recombinant vaccinia virus vaccine in man. *American Association of Cancer Research* (Abstract), (1994).
- 45 Correale, P., Zaremba, S., Nieroda, C., Zhu, M. Z., Schmitz, J., Schlom, J., and Tsang, K. Y. In vitro stimulation of human cytotoxic T lymphocytes specific for peptides derived from prostate specific antigen. *9th International Congress of Immunology* (Abstract), (1995).
- Disis, M. L., Smith, J. W., Murphy, A. A., Chen, W., and Cheever, M. A. In vitro generation of human cytolytic T-cells specific for peptides from the HER-2/neu protooncogene protein. *Cancer Research*, **54**:1071-1076, (1994).



Donovan, J. F., Lubaroff, D. M., and Williams, R. D. Immunotherapy of prostate cancer. *Problems in Urology*, 4:489-505, (1990).

5 Foon, K. A., Chakraborty, M., John, W., Sherratt, A., Kohler, H., and Bhattacharya-Chatterjee, M. Active immunity to the carcinoembryonic antigen (CEA) in patients treated with an anti-idiotypic monoclonal antibody vaccine. *Society for Biological Therapy* (Abstract), (1994).

10 Gauthier, E. R., Chapdelaine, P., Tremblay, R. R., and Dube, J. Y. Characterization of rhesus monkey prostate specific antigen cDNA. *Biochimica Biophysica Acta*, 1174:207-210, (1993).

15 Gritz, L., Destree, A., Cormier, N., Day, E., Stallard, V., Caiazzo, T., Mazzara, G., and Panicali, D. Generation of hybrid genes and proteins by vaccinia virus-mediated recombination: application to human immunodeficiency virus type 1 env. *J. Virol.* 64:5948-5957, (1990).

20 Helling, F., and Livingston, P. O. Ganglioside conjugate vaccines. Immunotherapy against tumors of neuroectodermal origin. *Molecular and Chemical Neuropathology*, 21:299-309, (1994).

25 Helling, F., Calves, M., Shang, Y., Oettgen, H. F., and Livingston, P. O. Construction of immunogenic GD3-conjugate vaccines. *Annals of the New York Academy of Science*, 690:396-397, (1993).

30 Huang, C. L., Brassil, D., Rozzell, M., Schellhammer, P. F., and Wright, G. L. Comparison of prostate secretory protein with prostate specific antigen and prostatic acid phosphatase as a serum biomarker for diagnosis and monitoring patients with prostate carcinoma. *Prostate*, 23:201-212, (1993).

35 Ioannides, C. G., Fisk, B., Fan, D., Biddison, W. E., Wharton, J. T., and O'Brian, C. Cytotoxic T cells isolated from ovarian malignant ascites recognize a peptide derived from the HER-2/neu proto-oncogene. *Cellular Immunology*, 151:225-234, (1993).

40 Irvine, K., Kantor, J., and Schlom, J. Comparison of a CEA-recombinant vaccinia virus, purified CEA, and an anti-idiotypic antibody bearing the image of a CEA epitope in the treatment and prevention of CEA-expressing tumors. *Vaccine Research*, 2:79-94, (1993).

45 Isaacs, J. T., Feitz, W. F., and Scheres, J. Establishment and characterization of seven Dunning rat prostatic cancer cell lines and their use in developing methods for predicting metastatic abilities of prostatic cancers. *Prostate*, 9:261-281, (1986).

Jenkins, S., Gritz, L., Fedor, C., O'Neil, E., Cohen, L. and Panicali, D. Formation of lentivirus particles in mammalian cells infected with recombinant fowlpox virus. *AIDS Research and Human Retroviruses* 7:991-998, (1991).

5

Kantor, J., Irvine, K., Abrams, S., Kaufman, H., Dipietro, J., and Schlom, J. Antitumor activity and immune responses induced by a recombinant carcinoembryonic antigen-vaccinia virus vaccine. *Journal of the National Cancer Institute*, 84:1084-1091, (1992a).

10

Kantor, J., Irvine, K., Abrams, S., Snoy, P., Olsen, R., Greiner, J., Kaufman, H., Eggensperger, D., and Schlom, J. Immunogenicity and safety of a recombinant vaccinia virus vaccine expressing the carcinoembryonic antigen gene in a nonhuman primate. *Cancer Research*, 52:6917-6925, (1992b).

15

Karr, J. F., Kantor, J. A., Hand, P. H., Eggensperger, D. L., and Schlom, J. Conservation of the prostate specific antigen (PSA) gene in primates and the expression of recombinant human PSA in a transfected murine cell line. *Cancer Research*:Submitted for Publication, (1995).

20

Kaufman, H., Schlom, J., and Kantor, J. A recombinant vaccinia virus expressing human carcinoembryonic antigen (CEA). *International Journal of Cancer*, 48:900-907, (1991).

25

Kleer, E., and Oesterling, J. E. PSA and staging of localized prostate cancer. *Urologic Clinics of North America*, 20:695-704, (1993).

30

Lilja, H. Structure, function, and regulation of the enzyme activity of prostate-specific antigen. *World Journal of Urology*, 11:188-191, (1993).

35

Livingston, P. O., Calves, M. J., Helling, F., Zollinger, W. D., Blake, M. S., and Lowell, G. H. GD3/proteosome vaccines induce consistent IgM antibodies against the ganglioside GD3. *Vaccine*, 12:1199-1204, (1993).

Lundwall, A., and Lilja, H. Molecular cloning of human prostate specific antigen cDNA. *FEBS Letters*, 214:317-322, (1987).

40

McEntee, M., Isaacs, W., and Smith, C. Adenocarcinoma of the canine prostate: immunohistochemical examination for secretory antigens. *Prostate*, 11:163-170, (1987).

Moss, B. Generation of recombinant vaccinia viruses. *Current Protocols in Molecular Biology*, 2:16.15.1-16.18.9, (1993).

45

Oesterling, J. E. Prostate specific antigen: a critical assessment of the most useful tumor marker for adenocarcinoma of the prostate. *Journal of Urology*, 145:907-923, (1991).

5 Panicali, D., Grzelecki, A. and Huang, C. Vaccinia virus vectors utilizing the  $\beta$ -galactosidase assay for rapid selection of recombinant viruses and measurement of gene expression. *Gene* 47:193-199, (1986).

10 Paoletti, E., Tartaglia, J., and Cox, W. I. Immunotherapeutic strategies for cancer using poxvirus vectors. *Annals of the New York Academy of Sciences*, 690:292-300, (1993).

15 Peace, D. J., Xue, B., Sosman, J. A., and Zhang, Y. In vitro immunization of human cytotoxic T lymphocytes specific for peptides derived from prostate specific antigen. *Cancer Vaccines: Structural Basis for Vaccine Development* (Abstract), (1994).

20 Powrie, F., and Coffman, R. L. Cytokine regulation of T-cell function: potential for therapeutic intervention. *Immunology Today*, 14:270-274, (1993).

25 Ravindranath, M. H., Brazeau, S. M., and Morton, D. L. Efficacy of tumor cell vaccine after incorporating monophosphoryl A (MPL) in tumor cell membranes containing tumor associated ganglioside. *Experimentia*, 50:648-653, (1994).

30 Ritter, G., Boosfeld, E., Adluri, R., Calves, M., Oettgen, H. F., Old, L. J., and Livingston, P. Antibody response to immunization with ganglioside GD3 and GD3 congeners (lactones, amide, and ganglisidol) in patients with malignant melanoma. *International Journal of Cancer*, 48:379-385, (1991).

35 Schellhammer, P. F., and Wright, G. L. Biomolecular and clinical characteristics of PSA and other candidate prostate tumor markers. *Urologic Clinics of North America*, 20:597-606, (1993).

40 Schlom, J., Kantor, J., Abrams, S., Tsang, K. Y., Panicali, D., and Hamilton, J. M. Strategies for the development of recombinant vaccines for the immunotherapy of breast cancer. *Breast Cancer Research and Treatment*, In Press.

45 Schroder, F. H. Experimental Models in the study of prostate cancer. Prostate Cancer. In: *International Perspectives in Urology*, 3:343-377, (1982).

Tsang, K. Y., Nieroda, C. A., De Filippi, R., Chung, Y. K., Yamaue, H., Greiner, J. W., and Schlom, J. Induction of human cytotoxic T cell lines directed against point-mutated p21 Ras-derived synthetic peptides. *Vaccine Research*, 3:183-193, (1994).

5

Wakui, S., Furusato, M., Nomura, Y., Asari, M., and Kano, Y. Lectin histochemical study of the prostate gland of the rhesus monkey (*Macaca mulatta*). *Journal of Anatomy*, 181:127-131, (1992).

10

Wang, M. C., Kuriyama, M., Papsidero, L. D., Loo, R. M., Valenzuela, L. A., Murphy, G. P., and Chu, T. M. Prostate antigen of human cancer patients. *Methods in Cancer Research*, 19:179-197, (1982).

15

Wang, M. C., Valenzuela, L. A., Murphy, G. P., and Chu, T. M. Purification of a human prostate specific antigen. *Investigations in Urology*, 17:159-163, (1979).

20

Zietman, A. L., Shipley, W. L., and Willett, C. G. Residual disease after radical surgery or radiation therapy for prostate cancer. Clinical significance and therapeutic implications. *Cancer*, 71:959-969, (1993).

## SEQUENCE LISTING

### (1) GENERAL INFORMATION:

- (i) APPLICANT: SCHLOM, JEFFREY  
PANICALI, DENNIS
- (ii) TITLE OF INVENTION: GENERATION OF IMMUNE RESPONSES TO  
PROSTATE-SPECIFIC ANTIGEN (PSA)
- (iii) NUMBER OF SEQUENCES: 2
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: SEWALL P. BRONSTEIN; DIKE, BRONSTEIN,  
ROBERTS & CUSHMAN
  - (B) STREET: 130 WATER STREET
  - (C) CITY: BOSTON
  - (D) STATE: MASSACHUSETTS
  - (E) COUNTRY: US
  - (F) ZIP: 02129
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: IBM PC compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
  - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER:
  - (B) FILING DATE:
  - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: RESNICK, DAVID S.
  - (B) REGISTRATION NUMBER: 34,235
  - (C) REFERENCE/DOCKET NUMBER: 44981
- (ix) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: (617) 523-3400
  - (B) TELEFAX: (617) 523-6440

### (2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 28 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: unknown
  - (D) TOPOLOGY: unknown

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

TCTAGAAGCC CCAAGCTTAC CACCTGCA

28

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 41 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: unknown

(D) TOPOLOGY: unknown

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

TCTAGATCAG GGGTTGGCCA CGATGGTGTC CTTGATCCAC T

41

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
21

This invention has been described in detail including the preferred embodiments thereof. However, it will be appreciated that those skilled in the art, upon consideration of this disclosure, may make modifications and improvements thereon without departing from the spirit and scope of the invention as set forth in the claims.

What is claimed is:

1. A method for generating an immune response to prostate-specific antigen (PSA) comprising, introducing a sufficient amount of a first pox virus vector to a host to stimulate an immune response, wherein the pox virus vector has at least one insertion site containing a DNA segment encoding PSA operably linked to a promoter capable of expression in the host.
2. The method of claim 1, further comprising at at least one periodic interval after introduction of the first pox virus vector contacting the host with additional PSA or a cytotoxic T-cell eliciting epitope thereof.
3. The method of claim 2, wherein the host is contacted with the additional PSA by introducing a second pox virus vector to the host having at least one insertion site containing a DNA segment encoding the PSA operably linked to a promoter capable of expression in the host.
4. A method for generating an immune response to prostate-specific antigen (PSA) in a host, comprising:
  - a. contacting the host with a sufficient amount of PSA or a cytotoxic T-cell eliciting epitope thereof; and
  - b. at least one periodic interval thereafter contacting the host with additional PSA or a cytotoxic T-cell eliciting epitope thereof.
5. The method of claim 4, wherein the host is contacted with the additional PSA by introducing a pox virus vector to the host having at least one insertion site containing a DNA segment encoding PSA or a cytotoxic T-cell eliciting epitope thereof operably linked to a promoter capable of expression in the host.



6. The method of claim 1 or 5, wherein the pox virus is selected from the group of pox viruses consisting of suipox, avipox, capripox and orthopox virus.

7. The method of claim 6, wherein the orthopox virus is vaccinia.

8. The method of claim 7, wherein the avipox is fowlpox, canary pox and pigeon pox.

9. The method of claim 8, wherein the suipox is swinepox.

10. The method of claim 3, wherein the first pox virus vector is vaccinia and the second pox virus vector is selected from the group of pox viruses consisting of suipox, avipox, capripox and orthopox virus.

11. The method of claim 2 or 4, wherein the PSA or T-cell eliciting epitope is formulated with an adjuvant or is in a liposomal formulation.

12. The method of claim 11, wherein the adjuvant is selected from the group consisting of RIBI Detox, QS21 and incomplete Freund's adjuvant.

13. A method for generating an immune response to PSA comprising contacting a host with a cytotoxic T-cell eliciting epitope of PSA.

14. The method of claim 13, wherein the T-cell eliciting epitope is formulated with an adjuvant or is in a liposomal formulation.

15. The method of claim 12, wherein the adjuvant is selected from the group consisting of RIBI Detox, QS21 and incomplete Freund's adjuvant.

16. A pharmaceutical composition comprising a pox virus vector having at least one insertion site containing a DNA reagent encoding PSA operably linked to a promoter and a pharmaceutical carrier.

## ABSTRACT

We have discovered that by using a recombinant viral vector, preferably a pox virus vector having at least one insertion site containing a DNA segment encoding prostate-specific antigen (PSA), operably linked to a promoter capable of expression in the host, a specific humoral and cellular immune response to PSA can be generated. The method preferably comprises introducing a sufficient amount of the recombinant pox virus vector into a host to stimulate the immune response, and contacting the host with additional PSA at periodic intervals thereafter. The additional PSA may be added by using a second pox virus vector from a different pox genus. In another embodiment, additional PSA can be added by contacting the host with PSA by a variety of other methods, including in one preferred embodiment adding PSA. The PSA may be formulated with an adjuvant or in a liposomal formulation.

1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398</
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	--------




Fig 1.

Fig 2A




Fig 2B





Fig 2C



DIKE, BRONSTEIN, ROBERTS & CUSHMAN  
130 Water Street  
Boston  
Massachusetts 02109

DECLARATION  
AND POWER OF ATTORNEY  
Original Application

ATTORNEY'S DOCKET NO. (If Any)  
45394

As a below named inventor, I hereby declare that

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed at 201 below) or an original, first and joint inventor (if plural names are listed at 201-203 below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

GENERATION OF IMMUNE RESPONSES TO PROSTATE SPECIFIC ANTIGEN (PSA)

which is described and claimed in:

☐ the attached specification or

☒ the specification in application Serial No. 08/500,306..... filed July 10, 1995.  
(for declaration not accompanying application)

And was amended on .....

if applicable

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, §1.56(a).

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed

FOREIGN APPLICATION(S), IF ANY, FILED WITHIN 12 MONTHS PRIOR TO THE FILING DATE OF THIS APPLICATION			
COUNTRY	APPLICATION	DATE OF FILING (day, month, year)	PRIORITY CLAIMED UNDER 35 U.S.C. 119
			YES NO
			YES NO
ALL FOREIGN APPLICATIONS, IF ANY, FILED MORE THAN 12 MONTHS PRIOR TO THE FILING DATE OF THIS APPLICATION			

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

(Application Serial No.)	(Filing Date)	(Status) (Patented, pending, abandoned)

**POWER OF ATTORNEY:** As a named inventor, I hereby appoint the following attorney(s) with full powers of association, substitution and revocation to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. (list name and registration number)

Sewall P. Bronstein (Reg. No. 16,919)  
Donald Brown (Reg. No. 20,845)  
David G. Conlin (Reg. No. 27,026)  
George W. Neuner (Reg. No. 26,964)

Ernest V. Linek (Reg. No. 29,822)  
Linda M. Buckley (Reg. No. 31,003)  
Ronald I. Eisenstein (Reg. No. 30,628)  
Henry D. Pahl, Jr. (Reg. No. 20,438)

Peter J. Manus (Reg. No. 26,766)  
David S. Resnick (Reg. No. 34,235)  
Peter F. Corless (Reg. No. 33,860)  
Kevin J. Fournier (Reg. No. 34,333)

Brian L. Michaelis (Reg. No. 34,221)  
Cara Z. Lowen (Reg. No. 38,227)  
John L. Welch (Reg. No. 28,129)  
Milton Mc.K. Oliver (Reg. No. 28,333)

SEND CORRESPONDENCE TO:  
Sewall P. Bronstein, Esq.  
Dike, Bronstein, Roberts & Cushman  
130 Water Street  
Boston, Massachusetts 02109

DIRECT TELEPHONE CALLS TO:  
(name and telephone number)

David S. Resnick  
617/523-3400

201	FULL NAME OF INVENTOR	LAST NAME <b>SCHLOM</b>	FIRST NAME <b>Jeffrey</b>	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY <b>Potomac</b>	STATE OR FOREIGN COUNTRY <b>Maryland</b>	COUNTRY OF CITIZENSHIP <b>U.S.A.</b>
	POST OFFICE ADDRESS	POST OFFICE ADDRESS <b>10301 Sorrel Ave.</b>	CITY <b>Potomac</b>	STATE OR COUNTRY <b>Maryland</b> ZIP CODE <b>20854</b>
202	FULL NAME OF INVENTOR	LAST NAME <b>PANICALI</b>	FIRST NAME <b>Dennis</b>	MIDDLE NAME <b>L.</b>
	RESIDENCE & CITIZENSHIP	CITY <b>Acton</b>	STATE OR FOREIGN COUNTRY <b>Massachusetts</b>	COUNTRY OF CITIZENSHIP <b>U.S.A.</b>
	POST OFFICE ADDRESS	POST OFFICE ADDRESS <b>114 Nonset Path</b>	CITY <b>Acton</b>	STATE OR COUNTRY <b>Massachusetts</b> ZIP CODE <b>01720</b>
203	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY ZIP CODE

I hereby further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

SIGNATURE OF INVENTOR 201	SIGNATURE OF INVENTOR 202	SIGNATURE OF INVENTOR 203
DATE	DATE <b>10-25-95</b>	DATE

U.S. DEPARTMENT OF COMMERCE  
PATENT AND TRADEMARK OFFICE

DIKE, BRONSTEIN, ROBERTS &  
CUSHMAN  
130 Water Street  
Boston  
Massachusetts 02109

DECLARATION  
AND POWER OF ATTORNEY  
Original Application

ATTORNEY'S DOCKET NO. (If Any)  
45394

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed at 201 below) or an original, first and joint inventor (if plural name is listed at 201 202 below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

GENERATION OF IMMUNE RESPONSES TO PROSTATE SPECIFIC ANTIGEN (PSA)

which is described and claimed in:

☐ the attached specification or

☒ the specification in application Serial No. 08/500,306 ..... filed July 10, 1995.  
(for declaration not accompanying application)

And was amended on .....

if applicable

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, §1.56(a).

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed

FOREIGN APPLICATION(S), IF ANY, FILED WITHIN 12 MONTHS PRIOR TO THE FILING DATE OF THIS APPLICATION			
COUNTRY	APPLICATION	DATE OF FILING (day, month, year)	PRIORITY CLAIMED UNDER 35 U.S.C. 119
			YES NO
			YES NO
ALL FOREIGN APPLICATIONS, IF ANY, FILED MORE THAN 12 MONTHS PRIOR TO THE FILING DATE OF THIS APPLICATION			

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application.

(Application Serial No.)	(Filing Date)	(Status) (Patented, pending, abandoned)

**POWER OF ATTORNEY:** As a named inventor, I hereby appoint the following attorney(s) with full powers of association, substitution and revocation to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. (list name and registration number)

Sewall P. Bronstein (Reg. No. 16,919)  
Donald Brown (Reg. No. 20,845)  
David G. Conlin (Reg. No. 27,026)  
George W. Neuner (Reg. No. 26,964)

Ernest V. Linek (Reg. No. 29,822)  
Linda M. Buckley (Reg. No. 31,003)  
Ronald I. Eikenstein (Reg. No. 30,628)  
Henry D. Pahl, Jr. (Reg. No. 20,438)

Peter J. Manus (Reg. No. 26,766)  
David S. Resnick (Reg. No. 34,235)  
Peter F. Corless (Reg. No. 33,860)  
Kevin J. Fournier (Reg. No. 34,333)

Brian L. Michaels (Reg. No. 34,221)  
Cara Z. Lowen (Reg. No. 38,227)  
John L. Welch (Reg. No. 28,129)  
Milton Mc.K. Oliver (Reg. No. 28,333)


SEND CORRESPONDENCE TO:  
Sewall P. Bronstein, Esq.  
Dike, Bronstein, Roberts & Cushman  
130 Water Street  
Boston, Massachusetts 02109

DIRECT TELEPHONE CALLS TO:  
(name and telephone number)

David S. Resnick  
617/523-3400

201	FULL NAME OF INVENTOR	LAST NAME <b>SCHLOM</b>	FIRST NAME <b>Jeffrey</b>	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY <b>Potomac</b>	STATE OR FOREIGN COUNTRY <b>Maryland</b>	COUNTRY OF CITIZENSHIP <b>U.S.A.</b>
	POST OFFICE ADDRESS	POST OFFICE ADDRESS <b>10301 Sorrel Ave.</b>	CITY <b>Potomac</b>	STATE OR COUNTRY <b>Maryland</b>
202	FULL NAME OF INVENTOR	LAST NAME <b>PANICALI</b>	FIRST NAME <b>Dennis</b>	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY <b>Acton</b>	STATE OR FOREIGN COUNTRY <b>Massachusetts</b>	COUNTRY OF CITIZENSHIP <b>U.S.A.</b>
	POST OFFICE ADDRESS	POST OFFICE ADDRESS <b>114 Nonset Path</b>	CITY <b>Acton</b>	STATE OR COUNTRY <b>Massachusetts</b>
203	FULL NAME OF INVENTOR	LAST NAME	FIRST NAME	MIDDLE NAME
	RESIDENCE & CITIZENSHIP	CITY	STATE OR FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP
	POST OFFICE ADDRESS	POST OFFICE ADDRESS	CITY	STATE OR COUNTRY

I hereby further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

SIGNATURE OF INVENTOR 201 	SIGNATURE OF INVENTOR 202	SIGNATURE OF INVENTOR 203
DATE <b>11/20/95</b>	DATE	DATE

U.S. DEPARTMENT OF COMMERCE  
PATENT AND TRADEMARK OFFICE

Practitioner's Docket No. 700953-45394-CPA-2

PATENT

JCS13 D.B. PTO  
09/693121  
10/20/00

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of: Schlom et al.  
Application No.: Continuation of 08/500,306  
Filed: July 10, 1995  
For: GENERATION OF IMMUNE RESPONSES TO PROSTATE-SPECIFIC  
ANTIGEN (PSA)


Assistant Commissioner for Patents  
Washington, D.C. 20231

CHANGE OF ATTORNEY'S ADDRESS IN APPLICATION

Please send all correspondence for this application as follows:

Ronald I. Eisenstein  
NIXON PEABODY LLP  
101 Federal Street  
Boston, MA 02110

Please direct telephone calls to:  
(617) 345-6054

  
SIGNATURE OF PRACTITIONER

Reg. No. 34,235  
Tel. No.: (617) 345-6057

David Resnick  
NIXON PEABODY LLP  
101 Federal Street  
Boston, MA 02110

CERTIFICATE OF MAILING/TRANSMISSION (37 C.F.R. 1.8(a))

I hereby certify that, on the date shown below, this correspondence is being:


MAILING

FACSIMILE

0 deposited with the United States Postal Service  
with sufficient postage as first class mail in an  
envelope addressed to the Assistant  
Commissioner for Patents, Washington, D.C.  
20231.

0 transmitted by facsimile to the Patent and  
Trademark Office.

Date: 10/20/00

  
Signature

Patricia Turner

(Change of Attorney's Address in Application, page 1 of 1)



## SEQUENCE LISTING

### (1) GENERAL INFORMATION:

(i) APPLICANT: SCHLOM, JEFFREY  
PANICALI, DENNIS

(ii) TITLE OF INVENTION: GENERATION OF IMMUNE RESPONSES TO  
PROSTATE-SPECIFIC ANTIGEN (PSA)

(iii) NUMBER OF SEQUENCES: 2

#### (iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: SEWALL P. BRONSTEIN; DIKE, BRONSTEIN,  
ROBERTS & CUSHMAN

(B) STREET: 130 WATER STREET

(C) CITY: BOSTON

(D) STATE: MASSACHUSETTS

(E) COUNTRY: US

(F) ZIP: 02129

#### (v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk

(B) COMPUTER: IBM PC compatible

(C) OPERATING SYSTEM: PC-DOS/MS-DOS

(D) SOFTWARE: PatentIn Release #1.0, Version #1.30

#### (vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:

(B) FILING DATE:

(C) CLASSIFICATION:

#### (viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: RESNICK, DAVID S.

(B) REGISTRATION NUMBER: 34,235

(C) REFERENCE/DOCKET NUMBER: 44981

#### (ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (617) 523-3400

(B) TELEFAX: (617) 523-6440

### (2) INFORMATION FOR SEQ ID NO:1:

#### (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: unknown

(D) TOPOLOGY: unknown

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

TCTAGAAGCC CCAAGCTTAC CACCTGCA

28

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 41 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: unknown

(D) TOPOLOGY: unknown

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

TCTAGATCAG GGGTTGGCCA CGATGGTGTC CTTGATCCAC T

41

CCCTGCA